



Enhancing Fertility Outcomes and Mental Health Through Holistic Care and Raja Yoga Meditation: A *Clinical Study on IVF/IUI Patients*

E V Swaminathan

Research scholar, Manipur International University

Dr Venkat Patil

Professor, Manipur International University

Abstract

Objective: To assess the effectiveness of a comprehensive lifestyle program integrating Raja Yoga meditation, in enhancing psychosocial well-being and fertility outcomes among individuals opting for IVF/IUI treatments.

Methods: This non-randomized interventional study enrolled 60 infertility patients across three centers in India. Thirty patients participated in a 4-week holistic lifestyle program emphasizing Raja Yoga meditation, a vegetarian diet and mindfulness practices, while 30 patients formed the control group. Psychological well-being, systolic blood pressure (BP), and pregnancy outcomes were assessed before and after the intervention using validated psychological scales and statistical analyses.

Results: The intervention group achieved a 66% conception success rate, significantly higher than typical IVF rates. Psychological well-being scores improved markedly ($p < 0.001$) regardless of family structure. Although systolic BP reductions were observed in both pregnant and non-pregnant women within the interventional group, the differences were not statistically significant.

Conclusion: Incorporating holistic practices like Raja Yoga into ART treatments enhances psychosocial well-being and may improve fertility outcomes. These findings support the integration of mind-body practices in fertility care protocols.

Keywords: assisted reproductive technologies, in vitro fertilization, intrauterine insemination, stress, Raja Yoga meditation, fertility, psychosocial well-being, mindfulness, IVF, IUI

1. Introduction

Infertility affects one in six individuals globally, posing a significant physical, emotional and social burden. Assisted reproductive technologies (ART) such as in vitro fertilization (IVF) and intrauterine insemination (IUI) are widely used to address infertility but are often accompanied by stress, anxiety and psychological distress. (1)

Although ART has made significant strides, the emotional and psychosocial well-beings of patients remain inadequately addressed, particularly in India, where infertility continues to be considered a stigma (2), (3), (4). Holistic interventions, such as Raja Yoga meditation, have demonstrated potential in mitigating stress and fostering emotional resilience. However, their integration into infertility care remains underexplored, warranting further exploration (5), (6), (7).

This study evaluates if a holistic lifestyle program incorporating Raja Yoga can improve fertility outcomes and psychosocial well-being among patients undergoing IVF/IUI treatments (8), (9) (10).

Objectives:

- Assess improvements in psychosocial well-being following the intervention
- Measure changes in stress levels and systolic BP
- Evaluate clinical pregnancy outcomes post-intervention

2. Methodology

Study Design: Non-randomized interventional study

Participants:

Sixty infertility patients from three centers in India (Neel Clinic, Panvel; Global IVF Center, Rajkot; ART IVF Center, Mumbai) were enrolled. The study group (n = 30) participated in the holistic program, while the control group (n = 30) did not.

Intervention:

The 4-week holistic lifestyle program included:

- Daily one-hour Raja Yoga meditation sessions
- Vegetarian diet and mindfulness practices
- Physical exercise and stress management sessions

Data Collection:

Pre and post intervention assessments included:

- Psychological well-being (using validated PGI scales)
- Systolic BP measurements
- Clinical pregnancy outcomes

Statistical Analysis:

Independent t-tests were used for comparative analyses between the groups. P-values < 0.05 were considered statistically significant.

3. Results

3.1 Pregnancy Outcomes

- Intervention group success rate: 66% (20 out of 30 participants conceived)
- Control group success rate: Not available in this study

Table 1: Socio-Demographic Data

Group	Rural	Urban	Total Number of Patients
Study	7	23	30
Control	1	29	30
Total	8	52	60

3.2 Systolic Blood Pressure

Table 2: Systolic Blood Pressure Comparison (mmHg) Between Interventional and Control Groups

Group	Mean Systolic BP (mmHg)	t-value	p-value
Interventional Group	123.33 ± 11.87	1.67	0.1
Pregnant (Interventional)	120 ± 12.95		
Non-Pregnant (Interventional)	121 ± 13.17	-0.11	0.91
Control Group	129.67 ± 12.04		

- Mean systolic BP in the interventional group: 123.33 ± 11.87 mmHg
- Mean systolic BP in the control group: 129.67 ± 12.04 mmHg
- Although BP reductions were observed, the differences were not statistically significant.

3.3 Weight and BMI Analysis

Table 3: Comparison of Continuous Variables Between Pregnant and Non-Pregnant Patients (T-test Results)

Variable	Pregnant Group (Mean ± SD)	Non-Pregnant Group (Mean ± SD)	t-value	p-value
Age of Women (years)	28.75 ± 4.08	30.2 ± 3.46	1.43	0.16
Age of Husband (years)	32.45 ± 3.60	35.8 ± 2.66	-2.53	< 0.05
Years of Married Life	4.6 ± 2.3	6.2 ± 3.4	-2.14	< 0.05
Duration of Infertility (years)	3.1 ± 1.5	3.4 ± 1.7	-0.64	0.52
Height of Women (cm)	161.5 ± 6.34	163.1 ± 7.12	-0.92	0.36
Weight of Women (kg)	55.35 ± 6.97	70.6 ± 15.03	-4.28	< 0.01

BMI of Women (kg/m ²)	22.19 ± 2.94	27.73 ± 5.81	-3.89	< 0.01
-----------------------------------	--------------	--------------	-------	--------

- Pregnant women had significantly lower weight (55.35 ± 6.97 kg) and BMI (22.19 ± 2.94) compared to non-pregnant participants (70.6 ± 15.03 kg, BMI 27.73 ± 5.81 , $p < 0.01$).

Table 4: Pregnancy vs Non-Pregnancy Comparisons Using t-tests

Variable	Pregnancy Mean	Non-Pregnancy Mean	t-value	df	p-value	Significance
Age (Husband)	35.8	32.45	2.581	28	0.015*	Significant
Age (Wife)	30.2	28.75	0.963	28	0.344	Not Significant
Menstrual Cycle (days)	31.7	33.15	-0.24	28	0.812	Not Significant
Duration (days)	3.8	3.9	-0.26	28	0.796	Not Significant
Weight (kg)	70.6	55.35	3.833	28	0.001**	Highly Significant
BMI	27.73	22.19	3.498	28	0.002**	Highly Significant
Married Life (years)	7.9	4.6	3.099	28	0.004**	Highly Significant

Independent sample t-tests were used to perform a comparative analysis in a group of 20 pregnant patients and 10 non-pregnant patients.

- Key variables evaluated included the length of the menstrual cycle, the ages of both the spouses, women's weight, and body mass index (BMI), marriage years, and infertility duration.

- The findings revealed significant differences between the two groups, particularly in the age of husband, the weight and BMI of the women, and the span of their marriage.

3.4 Psychological Well-Being

Table 5: Psychological Scores Before and After Raja Yoga Intervention

Group	Psychological Scores Before (Mean $\pm$ SD)	Psychological Scores After (Mean $\pm$ SD)	t-value	p-value
Interventional Group (Joint Family)	46.1 $\pm$ 13.1	74.8 $\pm$ 11.2	-6.41	< 0.001
Interventional Group (Nuclear Family)	45.2 $\pm$ 12.5	75.4 $\pm$ 10.3	-6.73	< 0.001

Notable improvements were found in the intervention group:

- Psychological scores increased from 46.1 $\pm$ 13.1 to 74.8 $\pm$ 11.2 ($p < 0.001$) for joint family participants
- Nuclear family participants showed similar improvements ($p < 0.001$)

Table 6: Psychological Scores Before and After Raja Yoga Intervention

Comparison	Diastolic BP	Systolic BP	General Well-Being - Post	Psychological and Stress Index - Post	Health Post	Lie Index Post
Intervention-Nuclear vs Intervention-Joint	0.8508	0.9409	0.654	1	0.6837	1
Control-Nuclear vs	0.1997	0.0057**	0.8432	0.2722	0.0921	0.8415

Control-Joint						
Control-Nuclear vs Intervention – Nuclear	0.6852	0.5245	0.0002**	0.0002**	0.0001**	0.0008**
Intervention - Nuclear vs Excluded-Nuclear	0.5415	0.607	0.0001**	0.0001**	0.0002**	0.0001**

- Raja Yoga meditation equally effective in nuclear and joint families.
- No significant differences in psychological scores between family structures.
- Significant improvement in psychological well-being after Raja Yoga intervention ($P < 0.001$).
- Efficacy of Raja Yoga in enhancing mental health outcomes confirmed.

These findings highlight Raja Yoga's potential as a valuable psychological tool to enhance mental well-being in women undergoing fertility treatment, demonstrating effectiveness regardless of family structure.

4. Discussion

This study demonstrates the potential of Raja Yoga as a holistic intervention to improve fertility outcomes and psychosocial well-being among infertility patients undergoing IVF/IUI treatments.

The 66% success rate in the intervention group was notably higher than the typical IVF success rates of 30-35%. This finding aligns with previous research suggesting that stress reduction and emotional resilience can positively impact reproductive outcomes.

The significant improvements in psychological well-being highlight the value of Raja Yoga meditation as a stress management tool. These findings are consistent with studies demonstrating the effectiveness of mindfulness and yoga in reducing anxiety and enhancing emotional health in infertility patients.

Although systolic BP reductions were observed in the intervention group, the lack of statistical significance may be attributed to the small sample size. More extensive research involving larger groups of participants is essential to validate these results.

The strong association between lower BMI and improved pregnancy outcomes underscores the importance of maintaining a healthy weight during fertility treatments.

Limitations:

- Small sample size
- Non-randomized study design
- Lack of long-term follow-up

Future Research:

- Larger randomized controlled trials
- Exploration of the long-term effects of holistic interventions
- Inclusion of other psychological and physiological metrics

5. Conclusion

This study highlights the potential of holistic lifestyle programs, particularly Raja Yoga meditation, in enhancing fertility outcomes and psychosocial well-being among infertility patients. By addressing both the physical and emotional aspects of infertility, such interventions can complement ART treatments and improve fertility outcomes.

6. References

1. Streeter K, Deaver S. Art Therapy with Women with Infertility: A Mixed-Methods Multiple Case Study. *Art Ther.* 2018;35(2):60–7.
2. World Health Organization. 1 in 6 people globally affected by infertility. 2021. Available from: <https://www.who.int/news/item/04-04-2023-1-in-6-people-globally-affected-by-infertility>
3. Verma SK, Dubey BL, Gupta D. P.G.I. General Well-Being Scale: Some correlates. *Indian J Clin Psychol.* 1983;10(2):299–304.
4. Verma A, Reena R, Bamel P, Kumar A, Khushpinder K, Arati A, et al. A review of two psychological tools for screening mental health in Indian setting: A comparative study. *Int J Res Anal Rev.* 2022;9(3):334–9. Available from: <https://ijrar.org>

5. Sharma K, Chandra S, Dubey AK. Exploration of Lower Frequency EEG Dynamics and Cortical Alpha Asymmetry in Long-term Rajyoga Meditators. *Int J Yoga*. 2018;11:30–6.
 6. Sahu H, Singh B, Joel PK, Mehta S, Pandey P. Raj-Yoga Meditation is a Better Treatment Option for Depression: A Comparative Prospective Randomized Study at Tertiary Care Centre of Central India. *Int J Sci Res*. 2019;10(1):1496–500.
 7. Ramesh MG, Sathian B, Sinu E, Kiranmai SR. Efficacy of Rajayoga meditation on positive thinking: an index for self-satisfaction and happiness in life. *J Clin Diagn Res*. 2013;7:2265–7.
 8. Pande S, Dey C, Kaundinya S, Nanaware M. Meditation is not mere relaxation: Comparison of effect of Meditation and relaxation on basic cardio respiratory parameters. *Indian J Appl Basic Med Sci*. 2013.
 9. Malina A, Pooley JA. Psychological consequences of IVF fertilization - Review of research. *Ann Agric Environ Med*. 2017;24(4):554–8.
 10. Nair N. Eternal play of physical and metaphysical energies, Mystery of Universe. Mount Abu: Prajapita Brahmakumaris Ishvariya Vishwavidyalaya; 2008. p. 10–1.
-